



**ASJO**

ISSN 3006-2543 (Online)

ISSN 1990-3863 (Print)

# **AL-SHIFA JOURNAL OF OPHTHALMOLOGY**

An Open Access, Peer Reviewed, Quarterly Journal of  
AL-SHIFA TRUST EYE HOSPITAL

**Vol. 20, No. 3, July – September 2024**

Indexed in

WHO Index Medicus (IMEMR)

Asian Digital Library (ADL)

Pak Medinet

Pakistan Medical and Dental Council IP/033

ISSN 3006-2543 (Online)  
ISSN 1990-3863 (Print)

A  
S  
J  
O

# Al-Shifa Journal of Ophthalmology

---

Vol. 20, No. 3, July – September 2024

---

**QUARTERLY PUBLISHED**

- **Editorial: Ethical Consideration in Research and Practice**
- **Biometry Formulas in High Myopes**
- **Anterior Lamellar Recession vs. Blepharoplasty**
- **Risks for Multiple Sessions of Retinal Photocoagulation**
- **Refractive Error in Healthy Infants of Nepal**
- **Central Corneal Thickness: Ultrasound vs. Optical Pachymetry**
- **Ocular Manifestations of Noonan Syndrome**

Abstracts available at <https://www.asjoalshifaeye.org> and <http://www.pakmedinet.com/ASJO>  
Manuscript submission through online platform [ejmanager.com](http://ejmanager.com)

---

**Indexed in Index Medicus -EMR**

---

**Recognized by Pakistan Medical & Dental Council – IP/033**

---

## Al-Shifa Journal of Ophthalmology

Editorial inquiries should be addressed to Prof. Dr. Tayyab Afghani, Department of Orbit and Oculoplastics, Al-Shifa Trust Eye Hospital, Jhelum Road Rawalpindi, Pakistan.  
Tel: 0092 51 5487821-25, Fax: 0092 51 5487827; Email: [aqrcpio@yahoo.com](mailto:aqrcpio@yahoo.com) ;  
Website: [www.asjoalshifaeye.org](http://www.asjoalshifaeye.org)

- Editorial: Ethical Considerations in Ophthalmic Research and Practice** 86  
Mahmood Ali
- Mean Errors From The Target Refraction at 1 Month After Phacoemulsification Surgery in High Myopes: A Comparison Of SRK/T, Haigis And Holladay 1** 88  
Shafaq Najmi, Badaruddin Athar Naem, Tehmina Nazir, Fariha Taimur, Zawar Ali Rathore
- Comparison of Anterior Lamellar Recession With and Without Blepharoplasty For Upper Eyelid Cicatricial Entropion** 93  
Asima Rafique, Muhammad Shaheer
- Obesity, Physical Inactivity, and Duration of Diabetes Mellitus as Risk Factors For Multiple Sessions of Retinal Photocoagulation** 101  
Muhammad Kamran Khalid, Muhammad Marwat, Muhammad Sharjeel, Muhammad Usman Awan, Uroosa Kanwal
- Refractive Error Among Healthy Infants in Tertiary Eye Care Centre of Nepal** 106  
Dr Govind Gurung, Krishna Kant Gupta, Harikant Sah
- Comparison of Central Corneal Thickness Using Ultrasound and Optical Pachymetry** 111  
Nashmia Jalil Malik, Muhammad Azam Khan, Irfan Aslam Khattak, Ayisha Shakeel, Huma Zainab, Maria Saleem
- Ocular Features of A Rare Case of Noonan Syndrome in A Pakistani Population** 119  
Murtaza Sameen Junejo

# Obesity, Physical Inactivity, and Duration of Diabetes Mellitus as Risk Factors for Multiple Sessions of Retinal Photocoagulation

Muhammad Kamran Khalid<sup>1</sup>, Muhammad Marwat<sup>2</sup>, Muhammad Sharjeel<sup>3</sup>, Muhammad Usman Awan<sup>3</sup>, Uroosa Kanwal<sup>3</sup>

## Abstract:

**Objectives:** Diabetic retinopathy (DR) is evolving as one of the leading causes of legal blindness worldwide. There is an immense need for the prevention of this potentially blinding disorder. Research has been going on to determine modifiable risk factors to decrease the progression of DR. More advanced cases of DR need pan-retinal photocoagulation (PRP) for the prevention of potentially blinding complications of DR.

Our study aimed to evaluate obesity, physical inactivity, and duration of DM as risk factors for multiple PRP sessions and the severity of DR.

**Methods:** This was a cross-sectional comparative study conducted at the Department of Ophthalmology, Gomal Medical College, Dera Ismail Khan, Pakistan from January 2021 to June 2021. The sample consisted of consecutive patients of pan-retinal photocoagulation (PRP) laser procedures during this period at the Eye Unit, DHQ Teaching Hospital, Dera Ismail Khan, Pakistan.

**Results:** A total of 168 patients undergoing laser procedures were included in the study. Out of these 104 (61.9%) were male and 64 (38.1%) were female. Obesity and duration of DM >5 years were found to be statistically significant ( $p < 0.05$ ) risk factors for multiple PRP sessions and severity of DR, whereas physical inactivity was not a statistically significant risk factor for multiple PRP sessions.

**Conclusion:** Obesity and duration of DM >5 years are significant risk factors for multiple PRP sessions in our setup. Efforts should be made to control all modifiable risk factors for the prevention of the sight-threatening complications of DR. *Al-Shifa Journal of Ophthalmology* 2024; 20(3): 101-105. © Al-Shifa Trust Eye Hospital, Rawalpindi, Pakistan.

- 
1. Gomal Medical College Dera Ismail Khan.
  2. Muhammad College of Medicine, Peshawar.
  3. Gomal Medical College Dera Ismail Khan.
- 

Originally Received: 05 May 2024

Revised: 27 May 2024

Accepted: 30 May 2024

## Correspondence to:

Muhammad Kamran Khalid  
Gomal Medical College, Dera Ismail Khan.

## Introduction:

Diabetic retinopathy (DR) is the leading cause of blindness in the working-age group globally<sup>1</sup>. International Diabetes Federation has recently released an estimated figure of 537 million diabetics worldwide and 33 million people are affected in Pakistan.

A review study on diabetic retinopathy has estimated that globally 35% of people with diabetes mellitus (DM) had some form of DR, 7% had proliferative diabetic retinopathy (PDR), 7% had diabetic macular edema (DME), and 10% were affected by the vision-threatening stages of diabetic retinopathy<sup>2</sup>.

Among other measures, intra-vitreous injections of anti-vascular endothelial growth factors (VEGFs) and Pan-retinal photocoagulation (PRP) procedures have been widely used for the treatment of these vision-threatening complications. These treatment strategies aim to halt further vascular proliferation and exudation on the retina. These modalities cannot revert the tissue damage already caused by the effects of metabolic disturbances of DM. So there is a significant need to prevent tissue damage by controlling the risk factors responsible.

Apart from good metabolic control<sup>3</sup>, the duration of DM, associated hypertension<sup>4</sup>, smoking, nephropathy, pregnancy, smoking, obesity<sup>5</sup>, and anemia have been associated with the progression of DR and its complications. However, relatively recent studies such as the Action in Diabetes and Vascular Disease (ADVANCE)<sup>3</sup> and the Action to Control Cardiovascular Risk in Diabetes (ACCORD-Eye)<sup>4</sup> have shown a limit to the risk reduction for DR that can be achieved with better glucose and BP management alone, respectively. Also, the evidence supporting the relationship between other modifiable risk factors and the severity of DR is inconclusive<sup>6-9</sup>. There remains a need for retinal photocoagulation for the prevention of vision-threatening complications of DR.

It can easily be understood that the more severe the DR, the more frequent the need for PRP sessions. So, the need for multiple PRP sessions is an indirect indicator of the severity of DR. As our study was conducted on patients being treated with laser PRP for severe DR, it is presumed that patients receiving multiple PRP sessions have more severe DR. Our objective was to determine whether Obesity, Physical Inactivity, and Duration of DM are risk factors for Multiple PRP Sessions in our location.

### **Materials and Methods:**

This was a cross-sectional comparative study conducted at the Department of

Ophthalmology, Gomal Medical College, Dera Ismail Khan, Pakistan from January 2021 to June 2021. The sample consisted of consecutive patients receiving Green laser photocoagulation during this period at the Eye Unit, DHQ Teaching Hospital Dera Ismail Khan, Pakistan. Approval from the ethical committee of Gomal Medical College, Dera Ismail Khan was taken before starting the study.

A total of 168 patients receiving Green laser procedures were included in the study using consecutive sampling during this period. Green laser procedures were performed with a mono-spot slit-lamp delivery system, Nidek GYC-1000, Japan in all patients under topical anesthesia using a wide-field Mainster PRP contact lens.

The patients were divided into those who received only one session of PRP (Single PRP Session) and those who received more than one (Multiple PRP Session). Obesity was defined as Body Mass Index (BMI) >30 kg/m<sup>2</sup> and less than 30 were taken as non-obese. Physical inactivity was defined as the absence of exercise with dedicated time and place. Based on the duration of DM, the subjects were divided into those having <5 years and those having ≥5 years duration from the onset of DM.

Patients with other ocular (including dense cataract, glaucoma, uveitis) or systemic (joint disease, end-stage kidney disease) comorbidities were excluded.

The sample was described by frequency and percentages using SPSS version 20 software. Gender and Age of the patient were the demographic variables and Obesity, Physical inactivity, and Duration of Diabetes were our clinical/research variables. The clinical variables were compared with the number of PRP sessions using the Chi-square test and a p-value <0.05 was taken to be statistically significant.

### **Results:**

A total of 168 patients were included in the study. Out of these 104 (61.9%) were male and 64 (38.1%) were female. Mean age of

the patients was  $56.83 \pm 12.5$  years. The frequency distribution of Obesity, Physical

Inactivity, Duration of DM, and PRP Sessions are shown in Tables 1.

*Table 1: Frequency distribution*

Obesity	Frequency	Percent
Obese	26	15.5%
Non-Obese	142	84.5%
Total	160	100%
Physical Inactivity	Frequency	Percent
Absent	26	15.5%
Present	142	84.5%
Total	160	100%
Duration of DM	Frequency	Percent
<5 years	36	21.4%
>5 years	132	78.6%
Total	160	100%
PRP Sessions	Frequency	Percent
Single PRP Session	88	52.4%
Multiple PRP Sessions	80	47.6%
Total	160	100%

A comparison between the research variables (Obesity, Physical Activity, and Duration of DM) and PRP sessions is shown in Tables 2 to 4 respectively.

*Table 2: Comparison of Obesity: PRP Sessions*

PRP Sessions	Obesity		Chi-Square	p-value
Single Session	Obese	Non-Obese	10.590	0.001
	6	82		
Multiple Session	20	60		

*Table 3: Comparison of Physical Inactivity: PRP Sessions*

PRP Sessions	Physical Inactivity		Chi-Square	p-value
Single Session	Absent	Present	1.034	0.309
	16	72		
Multiple Session	10	70		

*Table 4: Comparison of Duration of DM: PRP Sessions*

PRP Sessions	Duration of DM		Chi-Square	p-value
Single Session	<5 years	>5 years	17.598	0.000
	30	58		
Multiple Session	6	74		

This is evident from the above tables that Obesity ( $p=0.001$ ) and Duration of DM  $>5$  years ( $p=0.000$ ) are statistically significant ( $p<0.05$ ) risk factors for multiple PRP Sessions, whereas Physical Inactivity ( $p=0.309$ ) is not statistically significant ( $p>0.05$ ) risk factor for multiple PRP Sessions.

### **Discussion:**

It is evident from Table 2 that Obesity is a statistically significant risk factor for multiple PRP sessions and so for the severity of DR. Dirani et al had concluded that obese people were 6.5 times more likely to have PDR as compared to normal weight<sup>10</sup>. Also, they have shown that higher Body-mass index (BMI) was significantly associated with any DR ( $p=0.02$ ). Moreover, they have also shown that neck circumference ( $p=0.03$ ) and waist circumference ( $p=0.01$ ) were also significantly associated with any DR. In contrast Hwang et al had shown that higher BMI ( $p=0.001$ ), larger waist circumference ( $p=0.047$ ) and higher total body fat ( $p<0.001$ ) were significantly associated with lower risk of vision-threatening DR.

In our study, physical inactivity is not a statistically significant risk factor for multiple PRP sessions ( $p=0.309$ ) and so, neither for the severity of DR. AlQabandi et al had published an extensive review on this subject in which they have linked decreased sedentary times and more physical activity to the delayed onset and progression of DR and its severity<sup>11</sup>. They also added that physical activity provides both protective and anti-inflammatory effects on the retina. In our study duration of DM  $>5$  years is a statistically significant risk factor for multiple PRP sessions and so for severity of DR (Table:7). Similar results have been shown by Jenchitr W et al in their study at 10 and 20 years of DM. They have shown that for subjects having less than 10 years of DM, the prevalence of NPDR varied from 13.11% to 22.91% and PDR varied from 2.15% to 2.42%. Whereas subjects having up to 20 years of DM, the

prevalence of NPDR was up to 42.86% and PDR was up to 10.20%<sup>12</sup>. Niazi et al have shown the duration of DM as an independent risk factor for both severity and progression of DR (OR 5.7 for 5 to 10 years and 32.3 for more than 10 years in cases of NPDR and OR  $2 \times 10^6$  for 5 to 10 years and  $2 \times 10^8$  for more than 10 years in cases of PDR).

### **Conclusions:**

Obesity and duration of DM  $> 5$  years are significant risk factors for multiple sessions of PRP and so for severity of DR in our location. Control of modifiable risk factors as much as possible can decrease the risk of progression of DR and so for the need of multiple PRP sessions.

### **References:**

1. Klein BE. Overview of epidemiologic studies of diabetic retinopathy. *Ophthalmic Epidemiol* 2007;14:179–183.
2. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010;87:4–14.
3. Ismail-Beigi F, Craven T, Banerji MA, et al. Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet*. 2010;376:419–430.
4. Beulens JW, Patel A, Vingerling JR, et al. Effects of blood pressure lowering and intensive glucose control on the incidence and progression of retinopathy in patients with type 2 diabetes mellitus: a randomised controlled trial. *Diabetologia*. 2009;52:2027–2036.
5. Zhu W, Wu Y, Meng YF, Xing Q, Jian-Jun Tao JJ, Lu J. Association of obesity and risk of diabetic retinopathy in diabetes patients: A

- meta-analysis of prospective cohort studies. *Medicine* (2018) 97:32.
6. Henricsson M, Nystrom L, Blohme G, et al. The incidence of retinopathy 10 years after diagnosis in young adult people with diabetes: results from the nationwide population-based Diabetes Incidence Study in Sweden (DISS). *Diabetes Care*. 2003;26(2):349–354.
  7. The UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 1998;352(9131):854–865.
  8. Klein R, Klein BE, Moss SE. Is obesity related to microvascular and macrovascular complications in diabetes? The Wisconsin Epidemiologic Study of Diabetic Retinopathy. *Arch Intern Med*. 1997;157:650–656.
  9. Klein R, Klein BE, Moss SE, et al. The Wisconsin epidemiologic study of diabetic retinopathy, III: prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch Ophthalmol*. 1984;102:527–532.
  10. Dirani M, Xie J, Fenwick E, Benarous R, Rees G, Tien Yin Wong TY, Lamoureux EL. Are Obesity and Anthropometry Risk Factors for Diabetic Retinopathy? The Diabetes Management Project. *Invest Ophthalmol Vis Sci*. 2011;52:4416–4421.
  11. AlQabandi Y, Nandula S, Boddepalli C, et al. (August 21, 2022) Physical Activity Status and Diabetic Retinopathy: A Review. *Cureus* 14(8): e28238.
  12. Jenchitr W, Samaiporn S, Lertmeemongkolchai P, Chongwiriyannurak T, Anujaree P, Chayaboon D, Pohikamjorn A. Prevalence of Diabetic Retinopathy in Relation to Duration of Diabetes Mellitus in Community Hospitals of Lampang. *J Med Assoc Thai* 2004; 87(11): 1321-6.
  13. Niazi MK, Akram A, Naz MA, Awan S. Duration of Diabetes as a Significant Factor for Retinopathy. *Pak J Ophthalmol* 2010, Vol. 26 No. 4: 182-6.

**Authors Contribution**

Concept and Design: Muhammad Marwat  
Data Collection / Assembly: Muhammad Sharjeel  
Drafting Muhammad Usman Awan  
Statistical expertise: Uroosa Kanwal  
Critical Revision: Muhammad Kamran Khalid