



ISSN 3006-2543 (Online)

ISSN 1990-3863 (Print)

AL-SHIFA JOURNAL OF OPHTHALMOLOGY

An Open Access, Peer Reviewed, Quarterly Journal of
AL-SHIFA TRUST EYE HOSPITAL

Vol. 21, No. 2, April – June 2025

Indexed in

Index Medicus- EMR
Asian Digital Library
Pak Medinet

Recognized by

Higher Education Commission (HEC), Pakistan
College of Surgeon and Physician (CPSP), Pakistan
Pakistan Medical and Dental Council (PMDC) IP/033

ISSN 3006-2543 (Online)
ISSN 1990-3863(Print)

A
S
J
O

Al-Shifa Journal of Ophthalmology

Vol. 21, No. 2, April – June 2025

QUARTERLY PUBLISHED

- **Editorial: Intravitreal vs Sub-Tenon Triamcinolone in DME**
- **Vitamin D Levels and Myopia in Children**
- **Three-Point Local Anesthesia for Ex-DCR**
- **Intracameral Lidocaine in Sutureless Cataract Surgery**
- **Aqueous Misdirection in Post-Trabeculectomy**
- **Academic Performance in Medical Students with Eye Disease**
- **Corneal Epithelial Changes Post-PRK**
- **Risk Factors for PRP Sessions in Diabetic Retinopathy**
- **Retinal Inner Layer Disorganization in Diabetes**
- **Heparinised Irrigation and Inflammation in Phacoemulsification**

Abstracts available at <https://www.asjoalshifaeye.org> and <https://www.pakmedinet.com/ASJO>
Manuscript submission through online platform [ejmanager.com](https://www.ejmanager.com)

Indexed in Index Medicus -EMR, Asian Digital Library (ADL)
Recognized by Higher Education Commission (HEC), Pakistan
Recognized by College of Surgeon and Physician (CPSP), Pakistan
Recognized by Pakistan Medical and Dental Council (PMDC) IP/003

Al-Shifa Journal of Ophthalmology

Editorial inquiries should be addressed to Prof. Dr. Tayyab Afghani, Department of Orbit and Oculoplastics, Al-Shifa Trust Eye Hospital, Jhelum Road Rawalpindi, Pakistan.

Tel: 0092 51 5487821-25, Fax: 0092 51 5487827; Email: aqrcpio@yahoo.com ;

Website: www.asjoalshifaeye.org

Editorial: Gene Therapy in Ophthalmology: The Future of Inherited Retinal Diseases	62
Mehmona Asgher	
Comparative Efficacy Of Intravitreal versus Posterior Sub-Tenon Triamcinolone Acetonide Injections For Diabetic Macular Edema	63
Mahwish Shahid, Anum Nadir, Fauzia Naureen, Uzma Rehman, Alizay Gohar, Summaya Anjum	
Effective Way Of Local Anesthesia For External Dacryocystorhinostomy (Ex-Der); A Three Point Infiltrative Local Anesthesia	70
Piya Muhammad Musammat Rafi, Muhammad Rizwan Khan, Shehzad Manzoor, Muhammad Jahan Zaib Khan, Sarfraz Ahmad Mukhtar, Asif Manzoor	
Correlation of vitamin D levels with Myopia in Children: A Cross-Sectional Survey at a Tertiary Care Hospital in Rawalpindi	76
Saba Afzal Shaikh, Nazia Mushtaq, Nyla Gill, Khadija Mohammad, Ayyaz Hussain Awan, Ayesha Tariq	
Intracameral Lidocaine: A Safe and Effective Anesthetic Option for Manual Sutureless Cataract Surgery	82
Qaim Ali Khan, Muhammad Usman Arshad Qureshi, Rabia Sharif Bhatti, Aunaza Maqbool, Yasir Iqbal Malik	
Aqueous Misdirection Syndrome in Post-trabeculectomy Patients	89
Rima Khan, Yousaf Jamal Mahsood	
Histological/pathological Evaluation of Post Photo-Refractive-Keratectomy (PRK) Induced Changes in Corneal Epithelial Thickness and its Impact on Physiological Eye Functions	97
Raja Faisal Zulfiqar, Muhammad Yousuf Khoso, Tayyaba Kazmi, Sadia Sundus, Irfan Ul Akba, Raheela Adil	
Association of Age Groups, Gender, Smoking, and Hypertension with Pan-Retinal Photocoagulation Sessions in Diabetic Retinopathy Patients in the Population of D.I. Khan	108
Muhammad Shoaib Khan, Muhammad Kamran Khalid, Muhammad Sharjeel, Muhammad Abdullah, Hidayatullah Mahsud	
Disorganization of Retinal Inner layers in Diabetic Patients	113
Nimra Akram, Mazhar Ul Hassan	
Effect of Peroperative Use of Heparinised Irrigating Solution on Postoperative Inflammation in Phacoemulsification	121
Erum Nasir, Intisar Ul Haq Rana, Sana Nadeem, Asfandyar Asghar, Naila Obaid, Tehmina Nazi	

Correlation of vitamin D levels with Myopia in Children: A Cross-Sectional Survey at a Tertiary Care Hospital in Rawalpindi.

Saba Afzal Shaikh¹, Nazia Mushtaq², Nyla Gill², Khadija Mohammad¹, Ayyaz Hussain Awan¹, Ayesha Tariq³

Abstract:

Objectives: To assess the link between vitamin D levels and myopia in children.

Methods: From August 2023 to January 2024, a cross-sectional survey was conducted at Farooq Hospital in Rawalpindi. The trial included 100 children with vitamin D levels <10 ng/mL, ages 3–12, after obtaining parental consent. Children with a history of refractive problems, ocular illnesses, or systemic ailments were excluded from the study. Serum vitamin D levels were measured, and cycloplegic and noncycloplegic refractions were performed. SPSS version 23 was used to do the statistical analysis, and the student's t-test was used, and significance was indicated by $p < 0.05$.

Results: Of the 100 children, 38 (38%) had myopia. mean age was 7.04 ± 2.38 years. The sample consisted of 54 females (54%), and 46 boys (46%). Myopia and vitamin D deficiency had no significant association ($p = 0.115$). However, myopia was linked to older age ($p = 0.005$). It was more common in females than males (55.3%).

Conclusion: This study raises the possibility that a vitamin D deficit may not substantially raise the prevalence of childhood myopia. Further research must look at various genetic and environmental factors that affect the development of myopia. *Al-Shifa Journal of Ophthalmology* 2025; 21(2): 76-81. © Al-Shifa Trust Eye Hospital, Rawalpindi, Pakistan.

1. Akhtar Saeed Medical College, Rawalpindi
2. Nescom Hospital, Islamabad
3. Holy Family Hospital, Rawalpindi Medical University.

Originally Received: 05 Jan 2025

Revised: 28 Jan 2025

Accepted: 5 Feb 2025

Correspondence to:

Saba Afzal Shaikh
Akhtar Saeed Medical College, Rawalpindi
irfan.saba@yahoo.com

Introduction:

Myopia, also referred to as nearsightedness, is characterized by difficulty seeing distant objects clearly while maintaining close vision.¹ It is the most common cause of refractive error in the pediatric age group.² High myopia increases the risk of pathologic ocular changes such as cataracts, glaucoma, retinal detachment, and myopic macular degeneration, all of which can lead to blindness.^{3,4} The global myopia prevalence has increased from 28.3% in 2010 to 35.8 % in 2023 & this prevalence is expected to reach 39.80% by 2050.^{5,6} It is more than 2 times higher in East Asians than in white people. Various local studies revealed a prevalence of 19-41% among schoolchildren in Pakistan.⁵ With its potential to negatively affect quality of life and link to a higher risk of significant eye

disorders later in life, the rapidly rising frequency of myopia worldwide, especially among youngsters and most drastically in East Asia, has aroused serious public health concerns. Low vision puts the child and his or her family in danger of poverty and ill health. This finally leads to major concerns regarding national and international development. Low vision will undoubtedly imperil the achievement of two critical global indices of development, education, and health.⁵ Therefore, understanding the underlying causes of myopia is critical for developing effective management and prevention strategies.

According to recent studies, environmental and lifestyle factors such as excessive near work and extended screen time have a considerable role in the initiation and progression of myopia.^{2,6} The importance of vitamin D has gained increased recognition among these factors. Potential vision damage is one of the many health problems that have been connected to vitamin D insufficiency. There is growing evidence that low vitamin D levels may be linked to an increased risk of myopia development in children. Though the precise nature and extent of this association are still being investigated,^{7,8,9} few studies have suggested that vitamin D affects the axial length of the eye & vitamin D levels are inversely associated with axial length.¹⁰ Vitamin D is synthesised in the skin in response to ultraviolet B (UVB) radiation from sunlight, and deficiency is commonly related to decreased outdoor activity and limited sun exposure—both of which are linked to myopia.¹¹ Some studies have reported that sunlight exposure is more correlated with reducing axial length than vitamin D by affecting the release of dopamine in the retina.¹² Other factors have also been in research for myopia as atropine and defocus hypothesis.^{13,14}

Given the growing concern about vitamin D insufficiency and its possible effects on eye health, understanding the link between vitamin D levels and childhood myopia could help develop preventive strategies

and interventions, thus saving future generations from the most serious complication of blindness as it is the easiest way to control this growing epidemic in future. Despite extensive research in the West, the role of vitamin D in myopia development is still debatable, and various studies have shown controversial results. The authors were able to locate very few local studies on this crucial topic after a thorough review of the local literature. This cross-sectional survey seeks to look into the link between childhood myopia and vitamin D levels. By analyzing the association between these variables, this study aims to contribute to the better understanding of myopia's etiology and to influence future public health programs aimed at lowering the prevalence of this refractive error, as this is an easily preventable and treatable cause.

Methodology:

This cross-sectional survey was carried out at Farooq Hospital from August 2023 to January 2024. 100 children of both sexes, ages 3 to 12, who presented to paediatric OPD with vitamin D levels below 10 ng/ml were included in the trial after written parental agreement was obtained. The sample size was calculated using the formula $n = z^2 \cdot \text{lakP}(1-P)/d^2$ keeping a confidence level of 95%, an anticipated population proportion of 80, and an absolute precision of 0.08 (14). Children less than 3 or more than 12 years, already having any other refractive error, cataract, glaucoma, corneal/retinal disorders or congenital malformation of the eye, systemic disorders like Diabetes Mellitus, connective tissue disorders (Marfan & Ehler Danlos syndrome), Downs syndrome, and vitamin D and calcium supplements were excluded from the study. The ethical review board of Akhtar Saeed Medical College, Rawalpindi, reviewed & approved the whole protocol. An ophthalmologist from Farooq Hospital conducted the eye examination of the patients in Eye OPD. All patients were subjected to noncycloplegic

measurements of autorefractor for both eyes using Canon RK-F1 full autorefractor followed by cycloplegic refraction for the accurate diagnosis of myopia. Myopia was defined as ≤ -0.5 DS. A pre-designed questionnaire was used to gather data. The patient's serum vitamin D (1,25 dihydroxycholecalciferol) levels were measured using 2 millilitres of blood, which was then forwarded to the laboratory at Farooq Hospital. With SPSS version 23, statistical analysis was carried out. Welch Two sample t-test was applied. A p-value less than 0.05 was considered significant.

Results:

The analysis was performed for the 100 patients, who had a mean age of 7.04 ± 2.38 years. There were 54 girls (54%) and 46 boys (46%) having vitamin D deficiency. Myopia was seen in 38 patients, while 62 patients were found to have normal refractive errors. Of the 38 myopic patients, 21 (55.3%) were girls, while 17 (44.7%) were boys (Figure 1). The association of myopia with age is evident from Figure 2 ($p = 0.005$). Lastly, there was no significant association between myopia and vitamin D deficiency (Figure 3). Both groups had an almost equal range of vitamin D levels, as shown in Table 1.




Figure 1: Gender distribution of myopic children




Figure 2: Age distribution of myopic patients (years)




Figure 3: Association of vitamin D and Myopia

Table 1: Vitamin D levels in Myopic and normal children

	Vitamin D levels (ng/dl)	P value
Normal children	6.5-9	0.115
Myopic children	6.8-9.5	

Discussion:

In this study, the authors tried to find out the association of myopia and vitamin D levels in a pediatric population. There was no discernable association of myopia and vitamin D deficiency. With a sample of 100 children (mean age 7.04 ± 2.38 years), the findings showed that 38 patients (38%) were myopic, while 62 patients had normal vision. These results are in contrast to a local study that reported a mean age of 10 years with a predominance of boys, and myopia was seen in 55.6% of the children.¹⁵ Interestingly, 55.3% of the myopic youngsters in this study were female, which is consistent with findings from other studies that have shown a higher prevalence of myopia in females.¹⁶ According to another study, 51% of the participants were males, and the average age was 11.7 years.⁹ A study conducted in China reported the shift of age of myopia from 10 years in 2005 to 7 years in 2021.⁵

The lack of a significant relationship between vitamin D deficiency and myopia ($p = 0.115$) diverges from some existing literature that suggests a possible link between low vitamin D levels and an increased risk of myopia. For instance, a study by Tideman et al. (2016) reported that children with lower vitamin D levels had a higher prevalence of myopia.¹⁰ The same findings are reported by other studies as well.^{7,8,16} This lack of association could be because of several reasons such as limitations of the study design (such as cross-sectional data), confounding factors (such as outdoor activity, dietary intake of vitamin D or genetic predispositions), or insufficient power due to a small sample size. Moreover, there could be a threshold effect at play, where moderately low levels of vitamin D might have a different impact on myopia development compared to very

low levels. It is plausible that the association between vitamin D and myopia is stronger at a certain threshold of deficiency but may not be as pronounced in cases of extreme deficiency, where other factors may overshadow any potential effects of vitamin D on ocular development. However, the authors found few studies that support the findings of the current study with no association between low vitamin D levels and myopia.^{15,17-20} A study conducted in the UK found a significant relationship between myopia and fewer outdoor activities; however, they failed to find an association between low vitamin D levels and myopia, suggesting some unknown factor to be associated with outdoor activities that may have a protective effect.²² Another study conducted in 2021 revealed that sunlight exposure affects the development of myopia by affecting the release of dopamine in retina.¹² A systematic review conducted in 2023 reviewed various studies done to correlate the effects of nutrition, including vitamin D, with myopia. This review also failed to find any significant role of vitamin D in the development of myopia.²³ The current study supports the notion that other factors may play a more crucial role in the development of myopia, particularly age. The significant correlation found between older age and myopia ($p = 0.005$) corroborates findings by Araj et al. (2022), which indicated that the progression of myopia is significantly associated with the age of onset, higher weight, and body mass index.¹⁵ Similar results have been found in other studies that support the increasing prevalence of myopia with increasing age. It could be related to the increased growth rates causing a rapid increase in axial ocular length and environmental factors (increased

screen time and more near work due to the heavy academic burden and fewer outdoor activities).^{16,24}

The prevalence of myopia observed in our sample aligns with global trends. The rising incidence of myopia among children has been linked to lifestyle changes, including increased screen time and reduced outdoor activities.¹⁹ These environmental influences may overshadow the potential impact of vitamin D levels, suggesting that factors such as genetic predisposition and visual behaviour are more critical in myopia development.

Limitations of the current study include a small sample size and preselection of the patients with severe vitamin D deficiency, which may reduce the statistical power and generalisability of the findings to the broader population. The study does not account for other environmental factors (e.g., outdoor activity levels, screen time, seasonal variations) that may also influence myopia development and other potential confounding factors, such as genetic predispositions or nutritional status, which may affect the relationship between vitamin D and myopia. The absence of a significant association in the current research shows the necessity for further research as prospective control studies with larger sample sizes and various demographics to evaluate the relationship between vitamin D and myopia as well as comparison with controls with normal vitamin D levels.

Conclusion:

Findings of the current study indicate that Vitamin D may not be a significant risk factor in the development of myopia in children. Future research, especially longitudinal multicentre studies, should take a multifactorial approach, taking into account environmental, genetic, and behavioural factors that may contribute to the rising rates of myopia in the paediatric population.

References:

1. Limwattanayingyong J, Amornpetchsathaporn A, Chainakul M, Grzybowski A, Ruamviboonsuk P. The Association Between Environmental and Social Factors and Myopia: A Review of Evidence From COVID-19 Pandemic. Vol. 10, *Frontiers in Public Health*. Frontiers Media S.A.; 2022.
2. Kaur K, Gurnani B, Kannusamy V. Myopia: Current concepts and review of literature. *TNOA Journal of Ophthalmic Science and Research*. 2020;58(4).
3. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. *Ophthalmology*. 2016 May 1;123(5):1036–42.
4. Sankaridurg P, Tahhan N, Kandel H, Naduvilath T, Zou H, Frick KD, et al. IMI impact of myopia. *Invest Ophthalmol Vis Sci*. 2021;62(5).
5. Latif MZ, Khan MA, Afzal S, Gilani SA. Prevalence of refractive errors; an evidence from the public high schools of Lahore, Pakistan. *Journal of the Pakistan Medical Association* [Internet]. 2019 Apr;69(4):464–7. Available from: <https://www.researchgate.net/publication/332060879>
6. Huang HM, Chang DST, Wu PC. The association between near work activities and myopia in children - A systematic review and meta-analysis. *PLoS One*. 2015 Oct 20;10(10).
7. Veleva N, Dimtrova G, Oscar A, Kemilev P, Mladenov O, Dimitrov G, et al. Vitamin D status in children with myopia. *Bulgarian Review of Ophthalmology*. 2020;64(1).
8. Gao F, Li P, Liu YQ, Chen Y. Association study of the serum 25(OH)D concentration and myopia in Chinese children. *Medicine (United States)*. 2021;100(26).

9. Yu CY, Dong L, Li YF, Wei W Bin. Vitamin D and myopia: a review. *Int Ophthalmol*. 2024;44(1).
10. Tideman JWL, Polling JR, Voortman T, Jaddoe VWV, Uitterlinden AG, Hofman A, et al. Low serum vitamin D is associated with axial length and risk of myopia in young children. *Eur J Epidemiol*. 2016;31(5):491–9.
11. Al-Ajlan BY, Freije A, Allehdan S, Perna S. Prevalence and Risk Factors for Vitamin D Deficiency in Children and Adolescents in the Kingdom of Bahrain. *Nutrients*. 2023;15(3):494.
12. Landis EG, Park HN, Chrenek M, He L, Sidhu C, Chakraborty R, et al. Ambient light regulates retinal dopamine signaling and myopia susceptibility. *Invest Ophthalmol Vis Sci*. 2021 Jan 1;62(1):28.
13. Chia A, Lu QS, Tan D. Five-Year Clinical Trial on Atropine for the Treatment of Myopia 2 Myopia Control with Atropine 0.01% Eyedrops. *Ophthalmology*. 2016;123(2):391–9.
14. Fritschi A, Gerber C, Eggler D, Loertscher M. Simultaneous Myopic Defocus for Myopia Control: Effect on Accommodation, Peripheral Refraction and Retinal Image Quality in Non-Presbyopic Patients. *Optics*. 2021 Dec 1;2(4):200–15.
15. Aaraj S, Kausar A, Khan SA. Vitamin D deficiency: A risk factor for myopia in children — a cross sectional study in a tertiary care centre. *J Pak Med Assoc*. 2022;72(6):1075–9.
16. Holhos LB, Holhos TL, Coroi MC, Petricau C, Lazar L. Correlation between plasmatic vitamin D level and refractive status in children with disabilities. *Romanian Journal of Pediatrics*. 2021;70(2).
17. Li X, Lin H, Jiang L, Chen X, Chen J, Lu F. Low Serum Vitamin D Is Not Correlated With Myopia in Chinese Children and Adolescents. *Front Med (Lausanne)*. 2022;9.
18. Chou H Da, Yao TC, Huang YS, Huang CY, Yang ML, Sun MH, et al. Myopia in school-Aged children with preterm birth: The roles of time spent outdoors and serum vitamin D. *British Journal of Ophthalmology*. 2021;105(4).
19. Malik YI, Malik A, Shabbir R, Zaman A, Talib M. Role of Vitamin D in Near Sightedness. *Pakistan Journal of Ophthalmology*. 2020;36(2):136–40.
20. Wolf AT, Klawe J, Liu B, Ahmad S. Association Between Serum Vitamin D Levels and Myopia in the National Health and Nutrition Examination Survey (2001–2006). *Ophthalmic Epidemiol*. 2024;31(3):229–39.

Authors Contribution

Concept and Design: Nazia Mushtaq
 Data Collection / Assembly: Nyla Gill
 Drafting: Khadija Mohammad, Ayyaz Hussain Awan
 Statistical expertise: Ayesha Tariq
 Critical Revision: Saba Afzal Shaikh